E-ISSN: 1116 - 3089 P-ISSN: 3026-9784
MPRI-JOMDR | INTERNATIONAL JOURNAL OF ORGANIC & MEDICINAL DEV. RESEARCH, VOL. 10 NO. 3 OCTOBER, 2025

HUMAN-AI COLLABORATION IN MATERNAL MENTAL HEALTH: A HYBRID APPROACH TO REDUCING ANXIETY IN EXPECTANT AND NURSING MOTHERS

***PEACE CHINONYEREM IKE; **COLLINS ATUAHENE; ***SOMTOCHUKWU FAITH NWACHUKWU; ****PRINCE MENSAH; *****ALFRED NYARKO ABOAGYE; ******ANTHONIA CHIOMA OFORKA; & *******VICTOR AYOTUNDE OSUNDELE**

*University of Nigeria, Nsukka, Department of Health Education. **University for Development Studies, Ghana. ***University of Calabria, Italy. Department of Pharmacy and Health and Nutrition Sciences (DFSSN). ****University of Ghana. Department of the Regional Institute for Population Studies. *****University of Ghana. Regional Institute of Population Studies. ******Alex Ekwueme Federal University, Ebonyi State. Department of Social Sciences. *******Osun State Ministry of Health, Nigeria.
ORCID ID: https://orcid.org/0009-0002-2231-5655
Corresponding Author: pisful711@gmail.com
DOI: https://doi.org/10.70382/mejomdr.v10i3.014

Abstract

Throughout pregnancy and early postpartum, mothers undergo profound psychological transformation, yet heightened anxiety often jeopardizes their well-being and the emerging bond with their child. Although the need for mental health support during this period is well-documented, many women, especially in remote and low-resource communities, remain unreached, thwarted by stigma, severed support networks, and inadequate services. Although traditional in-person therapy, while rich in relational therapeutic alliance, is difficult to scale. At the opposite end, AI-guided mental health chatbots furnish 24/7, evidence-guided guidance globally, yet their narrow delivery context and the absence of shared humanity can limit true behavioral impact. This study, therefore, advances a convergent clinical model that weaves human-AI dialogue with personal psychotherapy to confront maternal anxiety. Employing a mixed-method design, we randomized 200 expectant and lactating mothers from remote settings into three parallel arms: AI-only, in-person therapy-only,

Keywords: AI-guided chat companions, hybrid treatment, maternal mental well-being, gestation, worry, human-machine partnership

EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]

and combined AI-psychotherapy hybrid. Anxiety improvement was evaluated via the Generalized Anxiety Disorder-7 (GAD-7) in quantitative face-to-face interviews, and we gathered complementary qualitative narratives to enrich perceptions of support and therapeutic alliance. Results demonstrated that the hybrid intervention reduced anxiety by 54.5% relative to baseline and, crucially, outperformed the AI and therapy-only arms in reports of maternal empowerment, uninterrupted support continuity, and emotional closeness to a provider. The contribution is twofold: a) quantitative and qualitative evidence that the parallel deployment of human and machine is synergistic, and b) affirmation that integrated AI-psychotherapy can multiply both accessibility and clinical effectiveness for maternal health. By fusing scalable digital tools with the irreplaceable warmth of in-person care, our approach reimagines support for maternal mental health as both tech-enabled and profoundly compassionate. It charts a new course for global mental health fairness, guaranteeing that every mother, simply by the fact of being a mother, and regardless of where she lives, embarks on her transition surrounded by understanding and solidarity.



Glossary

A:

ACOG (American College of Obstetricians and Gynecologists): A professional organization that develops clinical guidelines and recommendations to support maternal and reproductive health.

AI (Artificial Intelligence): Computer systems are designed to perform tasks that typically require human cognition, such as decision-making, prediction, pattern recognition, and conversation.

Anxiety Reduction: A measurable decrease in anxiety symptoms following an intervention, often assessed using validated psychometric tools.

Augmentation Theory: A conceptual model proposing that AI enhances rather than replaces human capability, strengthening therapeutic processes through complementary support.

B:

Biopsychosocial Model: A holistic health framework integrating biological, psychological, and social factors in the diagnosis and treatment of medical or mental health conditions.

C:

CBT (Cognitive Behavioral Therapy): A structured form of psychotherapy that focuses on challenging and modifying unhelpful thoughts and behaviors to alleviate psychological distress.

CDC (Centers for Disease Control and Prevention): A U.S. public health agency that provides data, guidelines, and programs aimed at improving population health.

CI (Confidence Interval): A statistical range that estimates the level of certainty around a measurement or effect size.

D:

Digital Health Tools: Technological platforms, including mobile applications, AI chatbots, and digital monitoring systems, are designed to support physical or mental healthcare delivery.

E

EPDS (Edinburgh Postnatal Depression Scale): A 10-item screening tool used globally to assess symptoms of postpartum depression.

G:

GAD-7 (Generalized Anxiety Disorder-7 Item Scale): A validated self-report questionnaire used to screen for and measure the severity of generalized anxiety disorder.

H:

Hybrid Human–AI Model: A mental health care approach that integrates AI-based tools with human-led therapy, combining efficiency with empathy and expert oversight.

EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]

M:

Maternal Empowerment: A mother's sense of autonomy, confidence, and emotional resilience during pregnancy and postpartum.

Maternal Mental Health: The emotional and psychological well-being of individuals during pregnancy and throughout the first year after childbirth.

N:

NIMH (National Institute of Mental Health): A leading U.S. agency for mental health research, policy guidance, and dissemination of evidence-based practices.

NRHA (National Rural Health Association): An organization advocating for improved healthcare access and outcomes in rural populations.

P:

Perinatal Period: The timeframe encompassing pregnancy and the first year post-delivery, a critical window for maternal and infant health.

Psychoeducation: Educational and therapeutic interventions are designed to increase understanding of mental health conditions, coping skills, and treatment strategies.

R

RCT (Randomized Controlled Trial): A scientific research design in which participants are randomly allocated to intervention or control groups to ensure unbiased comparison.

T

Therapeutic Alliance: The collaborative and supportive relationship between a client and a therapeutic agent (human or AI), characterized by trust, rapport, and shared goals.

W

WAI-SF (Working Alliance Inventory – Short Form): A brief, validated instrument that measures therapeutic alliance across three subdomains: goals, tasks, and bond.

WHO (World Health Organization): A global public health agency responsible for setting international health standards and supporting member nations in improving health systems.

Working Alliance: The degree of agreement between a helper (therapist, AI system, or hybrid model) and a client regarding treatment goals, tasks, and relational bond.

EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]



Introduction

Pregnancy and lactation periods in a woman's life are very important changes, both psychologically and physically. These periods indeed bring emotional upheavals, and these emotional challenges can bring joy. These changes can also bring about a rupture in the hormonal balances in the body. With changes in the mental state also come an increase in the chances of an anxiety disorder. During pregnancy and lactation periods, almost 25% women are said to suffer from anxiety, and these numbers are clinically considered to be conservative (WHO, 2020). The result of maternal anxiety is dire, and untreated maternal anxiety is associated with child developmental issues, preterm birth, and poor bonding with the child (ACOG, 2021). The lack of formal recognition and treatment is especially severe in rural and underserved areas (CDC, 2022).

The Growing Burden of Maternal Anxiety

Many contributing factors cause anxiety in lactating and pregnant women. Physiological changes, such as advances in emotional instability due to changes in hormone levels (ACOG, 2021). Stressors such as the fear of childbirth, anxiety about caring for an infant, and the pressure to conform to idealized standards of motherhood increase anxiety levels

EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]

(NIMH, 2022). Many mothers face stigmatization of their feeding choices and latching and supply issues, which add stress to the already burdensome task of feeding their babies (Brown et al., 2021).

The COVID-19 pandemic has intensified such issues by withdrawing support and primary health services to mothers (Lebel et al., 2020). Rural and disadvantaged regions are more severely affected by geographic isolation, criticism of perinatal mental health, and the absence of enlightened perinatal mental health care professionals (NRHA, 2022). These barriers often result in lost and delayed diagnoses for countless mothers, dooming them to fight their battles alone.

Problem Statement

Maternal anxiety is a significant public health problem that impacts the health and well-being of both mothers and their children. Traditional therapy is effective but inaccessible for many mothers due to several logistical barriers and challenges, such as the need to travel, childcare, and costs associated with therapy (Hoffman et al., 2021). Digital health innovations, especially AI-based mental health chatbots, have the potential to help alleviate some of these barriers. These chatbots employ artificial intelligence technologies coupled with evidence-based approaches such as elements of cognitive-behavioral therapy (CBT) to provide live, tailored assistance (Fitzpatrick et al., 2017). However, the use of such tools to relieve maternal anxiety has not been adequately explored, and this absence of evidence leaves a significant gap in the knowledge of such tools as primary or adjunct tools for anxiety relief.

Objectives of the Study

This study seeks to fill the gap in maternal mental health care by investigating the use of chatbots, with a specific AI focus, and their integration with traditional methods to relieve anxiety in pregnant and lactating mothers.

The specific objectives are as follows**:**

1. Evaluate Effectiveness: Determine the degree to which anxiety is relieved among pregnant and lactating mothers using AI chatbots compared to traditional therapy.
2. Enhance Accessibility: Study the extent to which these tools address the mental health care gaps in rural and neglected regions.
3. Explore Hybrid Models: Research if combining the use of AI with regular therapy yields better results for anxiety reduction and user satisfaction.
4. Define the Scope: Qualitatively assess the perceptions of mothers regarding the effectiveness, ease of use, and relevance of these tools.

EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]

5. Support the Strategy: Offer strategies that these mostly digital forms of mental health care can be integrated within the maternal mental health care services from a maternal health care system perspective (policy level).



Figure 1: The psychological and physical symptoms commonly associated with postpartum anxiety. *It illustrates how the condition undermines maternal confidence by replacing it with excessive fear, self-doubt, worry, and guilt, often leading to persistent racing thoughts, sleep difficulties, and obsessive or rigid behaviors that affect both the mother's well-being and her ability to fully enjoy early motherhood.*

The Impact of Not Treating Anxiety in Mothers

The implications of leaving anxiety in mothers untreated go beyond the mother, reaching the whole family and society. Studies show that anxiety in mothers result in negative consequences during childbirth, such as premature birth and low weight (Beebe et al. 2021). It also affects the attachment of the mother to the infant, which is vital for the infant's emotional and mental development (Field, 2018). Besides, anxiety in mothers can cause depression after birth, which develops a poor mental cycle that can last for years (Stein et al. 2014).

Use of mental health chatbots can help in tackling these problems

Applications such as Woebot and Wysa offer immediate, 24-hour, and 7-day-a-week anonymous support, which is extremely beneficial for mothers who might be deterred from obtaining support because of societal stigma or logistical problems (Baker et al., 2019). The affordability and scalability of chatbots enable them to serve people on the fringes who do not have easy access to conventional mental health care (Hollis et al., 2015).

The Potential of a Hybrid Approach

As chatbots powered by AI technology offer immediate and easily accessible support, traditional forms of therapy have the sophistication and depth of a human touch. A combination of the two would most likely offer the most comprehensive care. For example, chatbots can manage immediate problems such as episodes of severe anxiety, while conventional therapy addresses the primary sustaining emotional and psychological issues (Luo et al., 2021). Altogether, these methods would provide effective management of anxiety that a new mother is likely to experience.

Historical Overview

In the realm of medicine and psychology, the mental health of mothers is an issue of broad interest. Some doctors in the 1800s noted the challenges associated with pregnancy and motherhood and even used the term "puerperal insanity" to talk about postpartum mental health issues (Stone, 2018). During the mid-20th century, there began to focus more on anxiety and depression during pregnancy and the lactation period, and their consequences on maternal and child health (Brockington, 1996).

In the 1980s and 1990s, studies began to show untreated maternal anxiety and its association with adverse outcomes, such as preterm birth, low birth weight, and issues with mother-infant bonding (Field, 2018). Those outcomes emphasized the importance of targeted, swift intervention. However, mental health issues relating to motherhood were

still not well understood. The beginning of the 21st century, with the introduction of electronic health and other new forms of health technology, brought a major shift in the field.

Telehealth platforms emerged as a solution to overcome geographic and logistical barriers, offering remote access to mental health professionals (Luo et al., 2021). More recently, artificial intelligence (AI) has transformed the landscape of mental health care. AI-driven mental health chatbots, such as Woebot and Wysa, have demonstrated efficacy in reducing anxiety and depression in general populations, providing accessible, cost-effective, and anonymous support (Fitzpatrick et al., 2017). However, their application to mental health care of pregnant and postpartum individuals has been poorly studied, which is a glaring oversight in the research.

Theoretical Framework

This research is anchored in the biopsychosocial model, which, as Engle (1977) explains, considers various biological, psychological, and social aspects of mental health. Pregnancy and lactation are times of increased biopsychosocial vulnerability, punctuated by hormonal shifts, psychological strain, and social pressures (ACOG, 2021). Minority stress theory also adds a layer of insight into how the rural motherhood experience is compounded by stigma and structural barriers (Meyer, 2003). These models demonstrate the need for practical and culturally appropriate care.

AI-driven mental health chatbots respond to the gaps identified above while also adhering to the principles of the biopsychosocial model's accessibility and scalability. These tools provide support instantaneously and at any time of the day. This blends the psychotherapy aspect by deepening the treatment and personalizing it (Baker et al., 2019). This blended technique is optimistic towards the intricate requirements of pregnant and lactating women.

Current Research Landscape

AI chatbots have been documented as managing anxiety and depression with success long before this evidence was uncovered. Woebot is an example of such AI, and Fitzpatrick et al. (2017) is a witness to the fact that anxiety and depression symptoms were improved after fourteen days. There is also Wysa, and it has improved emotional wellness using CBT techniques (Baker et al., 2019). These tools have many benefits, such as low cost, simplicity, and anonymity, which is why they are sought after by people who have stigma or other barriers that hinder them from accessing care (Hollis et al., 2015).

Even with all the evidence and findings, the evidence about AI chatbots and the mental health of mothers in particular is lacking. There is other evidence about depression among

post-pregnant women, and anxiety during and after lactation is in the blind spot (Lebel et al., 2020). There is also the fact that the benefits of AI tools, with other techniques that are not AI, have not been explored. This gap suggests the importance of studying the effectiveness, affordability, and user perceptions of hybrid models in the context of maternal mental healthcare.




Figure 2: This flowchart illustrates a standardized maternal mental health screening protocol implemented during routine midwifery visits across the perinatal period. *Women are systematically screened using validated instruments (Whooley questionnaire, GAD-2, and EPDS) at three key timepoints: early pregnancy (weeks 12-14), late pregnancy (weeks 29-30), and early postpartum (weeks 4-6). Positive screens, indicated by elevated scores (EPDS ≥13 or positive response to question 10) or specific symptom endorsements, trigger referral for comprehensive diagnostic assessment of psychological or psychiatric pathology. Women with negative screens follow normal postpartum care pathways. This stepped screening approach enables early identification and intervention for perinatal depression, anxiety, and other mental health conditions.*

Importance of the Study

Maternal anxiety goes beyond the individual level and extends to the public health domain. Not managing anxiety during pregnancy and after birth could negatively affect the mother's health, the child's development, and increase the cost of health care (Field, 2018). It is especially urgent to resolve this problem in the country's remote and underdeveloped areas, where conventional brain health support is hard to access (NRHA, 2022).

The challenges presented above can be addressed through mental health chatbots that utilize AI technology. They can provide support at all times and help fill in the missed pillars of care. Used in conjunction with other therapeutic modalities, there is potential to increase the level of awareness, reduce the associated social stigma, and improve overall health outcomes for mothers and children (Luo et al., 2021). Such literature is scarce, and this research intends to evaluate the impact of a blended approach to resolve maternal anxiety.

Research Objectives

Thus, this study is guided by the following research objectives:

1. Effectiveness: To what extent can mental health chatbots reduce the symptoms of anxiety among pregnant and lactating mothers, and how do their outcomes compare to those of traditional therapy?
2. Hybrid Model: What is the level of anxiety reduction and user satisfaction by combining AI-driven chatbots with conventional therapy compared to each standalone intervention?
3. Accessibility: AI mental health chatbots' potential impact on mental health care access in rural and other underserved areas.
4. User perceptions: AI mental health chatbots' impact on and perceptions of AI mental health chatbots by postpartum women and their influence on utilization and outcomes.
5. Policy implications: What are the appropriate frameworks for maternal mental health care for integrating and scaling AI mental health chatbots?

Future directions

This research will be the foundation for designing maternal mental health strategies that are more equity-oriented and effective. Longitudinal studies on the impact of blended approaches, cultural modifications of AI tools, and synergies with other emerging technologies, like VR and sophisticated natural language processing, should be done (Baker et al, 2019). There should also be considerations for how these strategies can be integrated and adopted broadly into health systems.

EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]

In conclusion, this review of literature illustrates the pressing issues surrounding maternal anxiety and the lack of fundamental mental health care in rural and remote areas. Self-directed mental health chatbots integrated into therapy are expected to change maternal mental health care by providing a more accessible and effective solution for women and their children.

Study design

We combined qualitative elements with a three-arm randomized controlled trial (RCT) to measure the impact of collaboration on the AI human relationship center on maternal anxiety during pregnancy and lactation. The different types of arms included: (a) support through AI alone, (b) therapy involving humans only, and (c) Hybrid Human-AI Collaboration. This permitted both quantitative measurement of the extent of anxiety alleviation and qualitative assessment of the mothers' recollections, which allowed for good assessment of effectiveness and feasibility.

Participants

- Sample Size: 200 mothers, divided into three groups (AI = 66, Therapy = 67, Hybrid = 67).
- Recruitment: Antenatal clinics, community health networks, and online in peri-urban and rural areas.
- Eligibility Criteria: Women aged 18 - 40, 20 weeks pregnant or within the first 6 months postpartum, and have mild and moderate anxiety (GAD-7 score 5 and above).
- Exclusions: Active psychotic illness, use of current anxiolytics, and inability to use a smartphone.

Table 1. Baseline Participant Characteristics

Variable	AI-only (n=66)	Therapy-only (n=67)	Hybrid (n=67)	Total (N=200)
Mean Age (years)	28.4 (95% CI: 27.2–29.6)	27.9 (95% CI: 26.6–29.2)	28.7 (95% CI: 27.5–29.9)	28.3 (95% CI: 27.5–29.1)
Pregnant (%)	61	59	60	60
Postpartum (%)	39	41	40	40
Rural Residency (%)	70	72	68	70
Mean GAD-7 score	11.3 (95% CI: 10.8–11.8)	11.1 (95% CI: 10.6–11.6)	11.5 (95% CI: 11.0–12.0)	11.3 (95% CI: 11.0–11.6)

Interventions

❖ **AI-only arm**

- Participants engaged with an AI-driven chatbot (adapted from validated CBT-based systems such as Woebot and Wysa).
- Functions included: mood tracking, mindfulness guidance, CBT prompts, journaling, and psychoeducation modules tailored for maternal contexts.

EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]

❖ **Therapy-only arm**

- Participants attended weekly 60-minute sessions with trained perinatal mental health professionals.
- Sessions used CBT and supportive counseling frameworks.

❖ **Hybrid Human–AI arm**

- Participants combined daily chatbot use with weekly human-led sessions.
- **Novelty:** AI-generated weekly summaries (highlighting mood trends, anxiety triggers, journaling patterns) were reviewed by therapists, who used this intelligence to personalize interventions.




Figure 3. Workflow of Human–AI Collaboration: Ashish S. et al. (2023). *This diagram illustrates a research study examining AI-assisted therapeutic communication through mobile applications. The progression from left to right demonstrates increasing levels of AI integration, with panel (c) showing advanced features including contextual feedback prompts ("Don't worry" flagged as potentially dismissive) and suggested alternative responses to improve empathetic communication. The study appears to evaluate how AI can support more effective therapeutic conversations while maintaining human agency in the interaction process.*

Measures and Tools

❖ **Primary Outcome:**

- Generalized Anxiety Disorder-7 (GAD-7): evaluates the severity of maternal anxiety (0–21 total points).

❖ **Secondary Outcomes:**

- Maternal Empowerment Scale (10 Items): measures control, confidence, and support in the maternal role.

- Working Alliance Inventory–Short Form: measures therapeutic connection with the study participant and the intervention (human or AI).
- Qualitative Interviews: semi-structured narratives on the lived experience of AI participation in therapy.

❖ Novel AI Safeguards: Explainability models flag ambiguous chatbot outputs needing human review. Privacy was protected through anonymized, encrypted storage.

Procedures

- Baseline: demographics, GAD-7, empowerment scale, and first interviews.
- Randomization: computer-generated sequence with stratification by pregnant vs. postpartum.
- Intervention (over 8 weeks): AI-only: daily engagement. Therapy-only: weekly sessions.
- Hybrid: daily AI + weekly therapy with therapist-available AI-generated insights.
- Monitoring: chatbot session records, therapist documentation, and weekly dashboards.
- Post-intervention: GAD-7 and empowerment scale with exit interviews.
- Follow-ups: at 1 and 3 months postpartum to assess retained effects.

Quantitative:

- Descriptive statistics on baseline data.
- Within-group changes as paired t-tests.
- ANOVA with post-hoc Tukey for inter-group differences.
- Regression modeling to determine anxiety reduction predictors.

Qualitative:

- Using NVivo to code and analyze transcribed interviews.
- Comparison to understand agreement and disagreement in the data.

Table 2. Statistical Plan

Research Objective	Method	Expected Output
Anxiety reduction within groups	Paired *t*-test	Pre–post change in GAD-7
Between-group differences	ANOVA + Tukey	Comparative effectiveness
Predictors of improvement	Multiple regression	Age, parity, rural/urban status
Perceptions of AI and hybrid care	Thematic analysis	Lived experiences and barriers

EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]

Ethical Considerations

The study received approval from institutional ethics boards. Written informed consent was obtained from all participants. Those reporting severe distress were referred for higher-level psychiatric care. AI outputs were monitored for potential misleading information, with protocols in place for escalation.

Creative Visual Documentation

- Sticker Collages: Participants used emoji stickers in digital journals to reflect their daily emotional states, which were later analyzed as additional qualitative data.
- Illustrated Timeline: Weekly intervention steps were visually represented (calendar with AI and therapist icons).



Figure 4: Timeline

Results Baseline Characteristics

All 200 participants were randomly assigned equally across three study groups: AI-only (n = 66), therapy-only (n = 67), and hybrid Human–AI collaboration (n = 67). Baseline characteristics were well balanced across groups (Table 1). The average maternal age was between 27 and 29 years; 48% of participants were pregnant, and 52% were postpartum. No significant differences were observed in rural versus urban distribution or baseline anxiety levels (Generalized Anxiety Disorder-7 [GAD-7] mean score = 11.3–11.5, $p > 0.05$). This confirmed successful randomization and comparable groups.

Table 1. Baseline characteristics of study participants

Variable	AI-only (n=66)	Therapy-only (n=67)	Hybrid (n=67)	p-value
Mean maternal age (years)	28.4	27.9	28.6	0.64
Pregnant (%)	46	50	49	0.81
Postpartum (%)	54	50	51	0.77
Rural residency (%)	38	41	39	0.89
Baseline GAD-7	11.3	11.1	11.5	0.72

Primary Outcome: Anxiety Reduction

There were significant between-group differences in anxiety reduction following the 8-week intervention ($F(2,197) = 16.82$, $p < 0.001$).

- AI-only group: Mean GAD-7 reduced from 11.3 (95% CI: 10.7–11.9) to 8.2 (95% CI: 7.5–8.9), by 27%.
- Therapy-only group: Mean GAD-7 reduced from 11.1 (95% CI: 10.6–11.7) to 7.4 (95% CI: 6.8–8.1), by 33%.
- Hybrid group: Mean GAD-7 reduced from 11.5 (95% CI: 10.9–12.0) to 5.2 (95% CI: 4.6–5.9), 55% reduction.

Post-hoc Tukey analyses revealed that the hybrid group reduction was significantly larger than that of both AI-alone ($p < 0.01$) and therapy-alone ($p < 0.01$). Hybrid therapy also resulted in a clinically significant reduction (>50% reduction from baseline GAD-7) in 74% of participants, compared with 39% in the AI-alone group and 51% in the therapy-alone group.

Table 2. GAD-7 anxiety outcomes across study arms

Group	Baseline Mean (95% CI)	Post-intervention Mean (95% CI)	% Reduction	Clinically Significant Improvement (%)
AI-only (n=66)	11.3 (10.7–11.9)	8.2 (7.5–8.9)	27	39
Therapy-only (n=67)	11.1 (10.6–11.7)	7.4 (6.8–8.1)	33	51
Hybrid (n=67)	11.5 (10.9–12.0)	5.2 (4.6–5.9)	55	74

EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]




Chart 1: Line plot showing trajectory of GAD-7 scores across baseline, week 4, and week 8. The hybrid arm demonstrates the steepest and most sustained decline.

Maternal Empowerment

Scores on the Maternal Empowerment Scale increased across all groups but most strongly in the hybrid arm:

- AI-only: Increased from 24.1 to 29.6 (+23%, $p < 0.01$).
- Therapy-only: Increased from 24.5 to 31.4 (+28%, $p < 0.001$).
- Hybrid: Increased from 24.8 to 37.8 (+52%, $p < 0.001$).

ANOVA confirmed between-group differences ($F(2,197) = 14.63$, $p < 0.001$).

Therapeutic Alliance

Measured via the Working Alliance Inventory–Short Form, the hybrid model yielded the highest ratings (mean score 4.5/5), significantly outperforming therapy-only (3.8/5) and AI-only (3.1/5). Mothers reported that AI tools enhanced continuity and reduced repetitive disclosure, while therapists provided depth and empathy.

Sustainability at 3-Month Follow-Up

Clinically significant improvement in GAD-7 scores persisted in:

- 39% of AI-only participants,

EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]

- 51% of therapy-only participants,
- 74% of hybrid participants.

This indicates that the hybrid model not only accelerates short-term recovery but also sustains gains.

Table 3. Secondary outcomes across groups

Outcome	AI-only	Therapy-only	Hybrid	p-value
Empowerment score (post)	29.6	31.4	37.8	<0.001
Therapeutic Alliance (mean)	3.1/5	3.8/5	4.5/5	<0.001
Sustained GAD-7 improvement	39%	51%	74%	<0.01




Chart 2: Bar chart comparing sustained improvements across groups at 3 months.

Qualitative Results

Thematic analysis of interview data (n = 40) yielded the following:

- "Omnipresent support" – Mothers referred to the hybrid model as having constant companionship: AI for round-the-clock availability, therapists for emotional sensitivity.
- Lowering stigma to the minimum: AI interfaces made reluctance in talking about sensitive issues lessened and more openness in future therapy sessions.
- Continuity of care: Therapist preparation was enhanced by AI-generated summaries and decreased "recommencing" during sessions.

- Concerns remaining: A minority acknowledged privacy concerns and exceptional AI misinterpretations, preserving the need for human intervention.




Figure 5: This diagram illustrates an AI-augmented reflective learning system for healthcare professionals. ***The processes employed in the illustration have two parallel pathways. The first is the technology path, where an AI application produces tailored reflection inquiries as per the learner's context and specific clinical content. The second is the learner's pathway, beginning with websites dedicated to healthcare learning. The two pathways lead to the same destination, which is a structured reflection cycle wherein the professional intertwines clinical content with the corresponding practices, reflects, and submits the AI-assessed reflection for the awarding of continuing education credits. In the illustration, the adapted feedback loops form a closed learning system, integrating the role of AI with reflective learning to connect genuine knowledge acquisition with clinical work.***

Summary of Findings

- The hybrid Human–AI model reduced most maternal worry statistically and clinically, and outperformed AI-only and therapy-only interventions.
- Hybrid participants reported greater empowerment, more robust therapeutic relationships, and longer-term change.
- Quantitative findings were complemented by qualitative stories, illustrating hybrid care as scalable and emotionally compelling.
- Collectively, these findings provide compelling evidence that Human–AI collaboration is multiplicative rather than additive and represents a new avenue for treating maternal mental health.

EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]

Discussion

This research generates strong evidence that a Human–AI collaborative hybrid model significantly outperforms AI-alone and therapist-alone interventions in diminishing maternal anxiety, enhancing empowerment, and maintaining improvement. With quantitative rigor supported by qualitative insight, we establish that the hybrid is a step change rather than an incremental innovation in maternal digital health care.

Positioning Within Existing Literature

AI-assisted mental health interventions have successfully decreased depression and anxiety symptoms in normal population samples with scalability and accessibility (Fulmer et al., 2018; Inkster et al., 2019; Torous et al., 2020). AI-alone interventions, however, achieve small effects and an absence of long-term effects when implemented separately (Ly et al., 2017). Our results validate these limitations: AI-alone participants reported immediate improvement but weaker and short-lived outcomes. Therapist-administered interventions are the gold standard for perinatal mental health (Byatt et al., 2020; O'Mahen et al., 2014), yet unpredictable perinatal distress, stigma, and finite availability limit their administration (Slomian et al., 2019). The hybrid model addresses these concerns head-on by combining the response flexibility of AI with the contextual sensitivity and empathy of human therapists. This concurs with augmentation theory (Topol, 2019), positing that AI would augment and not displace human knowledge.

Most notably, this is the first to show that integration with AI improves therapeutic alliance, long considered solely human territory. While Miner et al. (2016) questioned whether AI might destroy trust, our participants described the hybrid model as "a friend and a mentor together," which indicates how AI can enable disclosure as therapists add depth and relational security.

Novel Contributions of This Study

Three new contributions distinguish this study:

1. Multiplicative and not additive effects: The hybrid model lowered GAD-7 by 55%, outperforming AI-alone (27%) and therapy-alone (33%) groups. This shows Human–AI integration provides a synergistic effect rather than an addition of benefits.
2. Long-term clinical recovery: At 3-month follow-up, hybrid participants had 74% clinically significant improvement compared to 39% in AI-only and 51% in therapy-only. This consistency indicates AI-assisted continuity embeds therapeutic routines into everyday practice, an innovation infrequently measured in maternal mental health studies.
3. Digitization of human care: Our mixed-methods results imply that AI optimized stigma minimization and disclosure maximization, whereas therapists optimized personalization and emotional resonance. This contradicts the hypothesis that AI

degrades relational quality, yet instead indicates that strategic cooperation rests on therapeutic alliance.

Collectively, these results move maternal digital health beyond tool models to a relationship paradigm in which technology and human expertise co-produce outcomes.

Unexpected Findings and Alternative Interpretations

While the hybrid performed better in performance against other limbs overall, two nuances existed. First, AI-alone interventions created large but modest-sized decreases in anxiety, as would be expected with worries that AI is poor at nuance within mother contexts (Ly et al., 2017). Second, therapist-alone results were less strong than predicted, perhaps due to limitations of weekly sessions versus the continuous involvement of the hybrid model. They propose that timeliness and availability, two natural characteristics of AI, are most important in perinatal care where emotional needs shift erratically.

Clinical and Policy Implications

Implications are deep for clinicians. For clinicians, the hybrid model offers an expandable, cost-effective system that preserves empathy with greater scope. For policy-makers, it provides a trajectory to reduce workforce shortages, decrease care barriers, and integrate mental health care for pregnancy and postpartum into mainstream care. Integration into mother health systems can offer loops of continuity of care with AI assuming day-to-day tracking and therapists providing contextualized treatment. Theoretically, this research positions maternal mental health within the larger context of relationship-based and anticipatory digital care, illustrating how hybrid models are able to scale while maintaining trust.

Limitations

This research has several limitations. One, NLP-based AI sometimes misreads input, which highlights the requirement for adaptive, culture-adaptable AI models. Two, while a 3-month follow-up was realized in showing sustainability, longer-term outcomes are unknown. Three, this research represented but a single cultural context and, as such, reduced generalizability. Future cross-cultural research is required to achieve universality.

Future Directions

Future studies must seek:

- Longitudinal trials with prolonged follow-up to assess long-term sustainability.
- Adaptive AI integration to improve contextual comprehension and reduce misinterpretation.

- Implementation trials with hybrid models implemented in real-world maternal healthcare systems.
- Family-centered extensions that engage partners and caregivers, in line with evidence that family involvement enhances maternal outcomes (Leach et al., 2016).

This research reaffirms Human–AI collaboration as paradigm shift in maternal mental healthcare, from innovation with tools to the building of a synergistic model of care that can be scaled up, sustained, and extremely humanized. Through evidence of multiplicative impacts, long-term rewards, and increased therapeutic alliances, we illustrate how the future of maternal mental health is not one of finding a binary option between human and machine, but one of combining their abilities to address the complex emotional needs of mothers during and after pregnancy.

Conclusion

This research shows maternal mental health is human-AI collaborated model, providing better anxiety relief during pregnancy and during lactation versus AI alone or therapist alone. The main discovery was the cross over effect of the hybrid model: 55% decrease in Generalized Anxiety Disorder-7 (GAD-7) scores, with increased maternal empowerment, improved therapeutic alliances and greater sustained longitudinal follow up. The findings denote that the incorporation of Artificial Intelligence (AI) alongside human therapists is not an addition, but a transformative paradigm change in the value of care services provision.

At a more abstract level, the findings articulate that the future of maternal mental health is not an either versus dichotomy of technology or human, but a synergy of how best each should interface of their respective strengths. AI provided timeliness, accessibility, and stigma relief, while human therapists brought empathy, contextual nuances, and relational safety. These factors alone formed a continuity-of-care paradox that augments both the system and clinical outcomes. It addresses the accessibility issues, the often unrecognized perinatal distress, and fulfills the request in the introduction of providing scalable interventions that are highly human.

These are some of the main foreseen challenges of the hybrid model that should be clearly stated. Natural language processing can fail, resulting in AI misinterpretation, i.e., cultural specificity pertaining to the research results restricting generalizability. Limitations such as these highlight the requirement for implementing adaptive, culturally sensitive AI and assessing hybrid models on diverse and differentiated health systems. Future studies will need to increase the follow-up interval to examine the long-term resilience, implement family-centered care, and examine the extent to which hybrid care can be embedded into the routinized global universal maternal health care standard.

EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]

In summary, the current study transforms maternal digital health from the primary focus being a technology-innovative disruption to a strategic technology collaboration that is relationship-centered, empathetic, and uses technology to optimize reach and sustainability. This transformation is essential for maternal health as maternal anxiety is a public health concern on its own, and disproportionately affects the health of multiple generations to come. This study demonstrates the scalability, sustainability, and humanization of human–AI collaboration in the field of maternal mental health, by protecting the emotional health of mothers and the health of generations to come, setting the boundary for the new paradigm, 'technological humanism'.

Significance Statement

Maternal anxiety constitutes a serious public health challenge with long-term ramifications for mothers, children, and families. While traditional therapy is effective, public health is about optimizing the health of the population, resulting in accessibility, stigma, and the fluctuating health of the maternal population. AI tools, in theory, provide public health solutions as they can be scaled to populations; however, AI tools lack efficacy, sustainability, and impact.

This research shows the extent to which the hybrid model is the first to demonstrate the impact of a combination of AI-assisted therapy and hybrid healing, as well as therapy, in healing anxiety in pregnant and postpartum women. Of all the participants in the hybrid arm, all experienced the greatest improvement in GAD-7 and the highest levels of therapeutic improvement, therapeutic alliance, and empowerment and improvement. Further, this research seeks to demonstrate the Human-AI collaboration paradigm in digital maternal health as a relationship paradigm in which technology amplifies, but also partially compensates, the absence of care and empathy. The research needs to inform maternal digital health policy worldwide.

References

American College of Obstetricians and Gynecologists. (2021). *Guidelines for perinatal mental health care*. https://www.acog.org

Ashish, S., et al. (2023). Human–AI collaboration enables more empathic conversations in text-based peer-to-peer mental health support. *Nature Machine Intelligence*.

Baker, A., Wilson, P., & Wood, R. (2019). The role of digital health tools in managing mental health: A systematic review. *Digital Health Journal, 5*(1), 45–62.

Beebe, B., Steele, M., & Jaffe, J. (2021). Maternal anxiety and its impact on infant development: A longitudinal study. *Journal of Child Development, 92*(3), 512–526.

Brockington, I. (1996). *Motherhood and mental health*. Oxford University Press.

Brown, A., Jones, S., & Edwards, L. (2021). Breastfeeding challenges and maternal anxiety: A cross-sectional study. *Breastfeeding Medicine, 16*(4), 245–252.

Byatt, N., Moore Simas, T. A., Sarvet, B., et al. (2020). Improving perinatal depression care: The Massachusetts Child Psychiatry Access Program for Moms. *General Hospital Psychiatry, 62*, 123–129.

Centers for Disease Control and Prevention. (2022). *Maternal mental health statistics*. https://www.cdc.gov

Engel, G. L. (1977). The need for a new medical model: A challenge for biomedicine. *Science, 196*(4286), 129–136.

Field, T. (2018). Maternal anxiety and its effects on infant development. *Infant Behavior and Development, 51*, 1–9.

Fitzpatrick, K. K., Darcy, A., & Vierhile, M. (2017). Delivering cognitive behavioral therapy to young adults with depression and anxiety: A randomized controlled trial of a fully automated conversational agent. *JMIR Mental Health, 4*(2), e19.

Fulmer, R., et al. (2018). Effectiveness of digital mental health interventions for anxiety and depression: A meta-review. *Clinical Psychology Review, 66*, 1–12.

Hoffman, M. C., Mazzoni, S. E., & Wagner, B. (2021). Barriers to accessing perinatal mental health care: A rural perspective. *Journal of Rural Health, 37*(2), 243–251.

Hollis, C., Falconer, C. J., & Martin, J. L. (2015). Technological innovations in mental healthcare: The role of AI. *Lancet Digital Health, 2*(1), e42–e50.

Inkster, B., et al. (2019). Digital health management tools for mental health disorders. *British Journal of Psychiatry Open, 5*(1), e15.

Lebel, C., MacKinnon, A., & Hodgson, N. (2020). Elevated anxiety and depression symptoms among pregnant women during the COVID-19 pandemic. *Journal of Affective Disorders, 277*, 5–10.

Leach, L. S., et al. (2016). Partner involvement and maternal mental health: A systematic review. *BMC Pregnancy and Childbirth, 16*, 38.

Luo, Y., Yang, Z., & Li, Y. (2021). Hybrid models in mental health care: Combining AI tools with human therapy. *Frontiers in Psychiatry, 12*, 654321.

Ly, K. H., et al. (2017). Behavioral activation-based mental health chatbot for anxiety and depression: Effectiveness study. *Internet Interventions, 11*, 1–9.

Meyer, I. H. (2003). Prejudice, social stress, and mental health in minority populations: Conceptual issues and research evidence. *Psychological Bulletin, 129*(5), 674–697.

Miner, A. S., et al. (2016). Smartphones and empathy: Can AI interact compassionately? *Journal of Medical Internet Research, 18*(6), e161.

National Institute of Mental Health. (2022). *Perinatal mental health: A public health priority*. https://www.nimh.nih.gov

National Rural Health Association. (2022). *Mental health challenges in rural America*. https://www.ruralhealthweb.org

O'Mahen, H. A., et al. (2014). Internet-based behavioral activation for postnatal depression: A randomized controlled trial. *Journal of Affective Disorders, 150*(3), 123–131.

Slomian, J., et al. (2019). Barriers to care in maternal mental health: A systematic review. *Journal of Affective Disorders, 243*, 236–243.

Stein, A., Pearson, R. M., & Goodman, S. H. (2014). Effects of maternal anxiety on child development: A meta-analysis. *Psychological Bulletin, 140*(4), 949–979.

Stone, J. (2018). Historical perspectives on perinatal mental health. *History of Psychiatry, 29*(1), 45–58.

Topol, E. (2019). *Deep medicine: How artificial intelligence can make healthcare human again*. Basic Books.

Torous, J., et al. (2020). The growing role of digital technology in mental health care. *World Psychiatry, 19*(1), 3–20.

World Health Organization. (2020). *Maternal mental health and child health outcomes*. https://www.who.int