MPRI-JOMDR | INTERNATIONAL JOURNAL OF ORGANIC AND MEDICINAL DEVELOPMENT RES.
VOL. 1 NO. 1 JUN-2022

SYNTHESES, CHARACTERIZATION AND DETERMINATION OF ANTI-MICROBIAL ACTIVITIES OF 1-{(E)-[2-{(2-HYDROXYNAPHTHALENE-1-YL) METHYLIDENE] AMINO} ETHYL} NAPHTHALENE-2-OL) DERIVED FROM ETHYLENEDIAMMINE AND 2-HYDROXY-1-NAPHTHALDEHYDE AND ITS COMPLEXES WITH COBALT (II) AND NIKEL (II) METAL IONS

BULUS, CALEB LARAPS; DAUDA, ABUBAKAR YAKUBU; AND ELISHA, VICTOR ZUYA

Department of Science Laboratory Technology (SLT), Federal Polythecnic Bauchi.

Abstract

The Schiff base 1-{(E)-[2-{(2-hydroxynaphthalene-1-yl) methylidene] amino} ethyl} naphthalene-2-ol was derived from ethylenediamine and 2-hydroxy-1-naphthaldehyde and its complexes with nickel(II) and cobalt(II). The UV-Visible result showed molar absorptivity above 1000 $Lmol^{-1}cm^{-1}$ ($\pi \sim \pi^*$) in the complexes, the FT-IR result showed absorption about 1600 nm (N=C) in all. The ligand and the complexes were screen for anti-microbial activity and the cobalt (II) complex was found to inhibit growth in *B. subtilis*, *S. aureus*, *E. coli*, *S. sp* and *P. auruginosa* in that order, with minimum inhibition concentration in *B. subtilis at* 20 µg/ml which is comparable to standard drugs while

KEYWORDS:
synthesis, characterization, ethylenediammine , 2-hydroxy-1-naphthaldehyde, complexes, cobalt, nikel.

EDITOR-IN-CHIEF
Prof. A.I. Mteiye [MALAWI]

500 µg/ml in the others which is too high. The nickel (II) and the Schiff base ligand only have growth inhibition against *B. subtilis, E. coli,* and *S. sp* with minimum inhibition concentration at 500 µg/ml which is too high compared to standard drug values.

INTRODUCTION


A Schiff base (named after Hugo Schiff, a German Chemist) is a compound with the general structure $R_2C=NR^1$ ($R^1 \neq H$). They can be considered as sub-class of imines, being either secondary ketimines or secondary aldimines depending on their structure. The term is often synonymous with azomethine which refers specially to secondary aldimines.

The term Schiff bases is normally applies to these compounds when they are being used as ligands to form coordination complexes with metal ions. Such complexes do occur naturally, for instance in corrin, but the majority of Schiff bases are synthesized for a wide range of application (IUPAC, 1997). For instance, metals neodymium(III) and thallium(III) complexes of the Schiff base derived 4-chlorobenzenesulphonyl chloride and aminobenzothiazole were screened *in vitro* for their anti-bacterial activities against *E.coli, S.aureus* and *P.aeruginosa* and were found to be more active against the microorganisms screened relative to ciprofloxacin, gentamicin and co-trimoxazole (Obasi *et al*., 2017). Metal complexes of Cu^{2+}, Co^{2+}, Ni^{2+} and Mn^{2+} with Schiff bases derived from O-phthalaldehyde and amino acids such as glycine, L-alanine, and phenyl-alanine were tested against three fungi. It was found that Cu^{2+} and Ni^{2+} complexes exhibited growth inhibition towards the entire studied fungi. However Co^{2+} and Mn^{2+} complexes exhibited less growth inhibition (Neelakantan *et al*., 2008). Glutamic acid-salicylaldehyde Schiff base metal complexes were bound into bovine serum albumin (BSA), which afforded BSA binding Schiff base metal complexes (BSA-sal, GluM, M=Cu, Co, Ni, Zn). It showed the antioxidant capacity of BSA increased more than 10 times after binding Schiff base metal complexes (Wang *et al*., 2007). Schiff base derived from 4-aminoantipyrine and benzaldehyde derivatives were tested for anti-inflammatory activity and the result was promising which could be beneficial for use in the treatment of inflammatory diseases (Alam, *et al*., 2012).

PROCEDURE/METHOD

Synthesis of 1-{(E)-[2-{(2-hydroxynaphthalene-1-yl) methylidene] amino} ethyl} naphthalene-2-ol

The synthesis as carried out by Cleiton *et al.*, (2009). About 1ml (1.0g, 16.67 mol) of ethylenediammine was measured and dissolved in little amount of water, then 5.73g (33.31mol) of 2 – hydroxyl – 1 – naphthaldehyde was weighed and dissolved in methanol. The two solutions were mixed and refluxed while stirring with a magnetic stirrer at a temperature of 60 °C for about 30 min. A yellow coloured precipitate was formed which was filtered and dried in a desiccators. The dried solid product was recrystallized from DMSO by dissolving it in DMSO at high temperature and the resulting clear solution filtered while hot and allowed to cool. After cooling the yellow precipitate was reformed which was filtered then washed with a little quantity of methanol and dried in a desiccators, then weighed and labeled EDNA.



Ethylenediamine

2- hydroxy-1-naphthaldehyde

1-{(*E*)-[(2-{[(2-hydroxynaphthalen-1-yl) methylidene]amino}ethyl)imino]methyl} naphthalen-2-ol

Equation:5 Synthesis of Schiff base derived from ethylenediamine and 2-Hydroxy-1-naphthaldehyde

Complexations

The bidentate ligand 1-{(E)-[2-{(2-hydroxynaphthalene-1-yl) methylidene] amino} ethyl} naphthalene-2-ol was used to form chelate complexes with Ni (II) and Co (II) ions (Shetye, *et al*., 1999).

Nickel-ligand complexation

As use by Selma Bal and Sedat Salih Bal, (2014). Exactly 2.56g (20 mmol) of the $CoCl_2.6H_2O$ salt was weighed and dissolved with a minimum amount of water in a beaker and 3.68 g (10 mmol) of the ligand 1-{(E)-[2-{(2-

hydroxynaphthalene-1-yl)methylidene]amino}ethyl}naphthalene-2-ol was weighed and dissolved in DMSO on heating. The two solutions were mixed and heated with continuous stirring using magnetic stirrer for 30 min. A greenish-brown precipitate was formed which was filtered and dried in a desiccators. The dried precipitate was recrystallized from DMF by dissolving it at high temperature, filtered and allowed to cool. On cooling the precipitate crystallized out of solution and was filtered and dried in a desiccators then weighed and labeled NiEDNA and stored.

Cobalt-Ligand Complexation

As used by Arumugam et al (2017). Exactly 2.56g (20 mmol) of $CoCl_2.6H2O$ salt was weighed and dissolved with a minimum amount of methanol in a beaker and 68 g (10 mmol) of the ligand 1-{(E)-[2-{(2-hydroxynaphthalene-1-yl)methylidene]amino}ethyl}naphthalene-2-ol was weighed and dissolved in DMSO on heating. The two solutions were mixed and heated with continuous stirring using magnetic stirrer for 30 min. A reddish-pink precipitate was formed which was filtered and dried in a desiccators. The dried precipitate was recrystallized from DMF by dissolving it at high temperature, filtered and allowed to cool. On cooling the precipitate crystallized out of solution and was filtered and dried in a desiccators then weighed and labeled CoEDNA and stored.

Determination of minimum inhibitory concentration (MIC) of the complexes using the broth micro dilution method.

This was conducted at the laboratory of the department of pharmacy of the University of Nigeria, Nsukka.

1. From the stock (2500 µg/ml) of the complexes, five different concentrations (500, 100, 20, 4, 2 µg/ml) were prepared in Muellarhinton broth.
2. Fresh cultures of the test pathogens (*Bacillus subtilis, Staphylococcus aureus, Streptococcus sp., Escherichia coli* and *Pseudomonas aeruginosa*) were obtained and adjusted to a clear or less turbid suspension.

EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]

3. Exactly 100 µL of the adjusted cell suspension is introduced into the Muellarhinton broth containing different concentrations of the complexes.
4. This is followed by a 24 h-incubation at room temperature.
5. The minimum inhibitory concentration is defined as the lowest concentrations of the complexes which prevented visible growth of the test pathogen.

Spectroscopic Analyses

FT – IR (Fourier Transformed Infrared)

The FT – IR Spectroscopic analysis was carried out at Agilent Technologies in Zaria Kaduna State, Nigeria with IR-1900 series and the result reported in Chapter Four of this work.

UV-Visible Spectroscopic analysis of the samples

A UV-1800 series spectrophotometer was used to determine the absorbance and wavelength of absorptions in the range of 200 nm-700 nm of the samples dissolved in DMSO in a 1 cm cell with sample concentration of 0.0005 M and the result reported in chapter four.

Result, Discussion and Conclusion

Results

Percentage yield of:

(i) The 1-{(E)-[2-{(2-hydroxynaphthalene-1-yl) methylidene] amino} ethyl} naphthalene-2-ol ligand

$$\% \text{ yield} = \frac{\text{Actual yield}}{\text{Theoretical yield}} \times 100\%$$

Actual yield = 5.41g

Theoretical yield = 6.13g

% yield = 88.25%

(ii) Ni-1-{(E)-[2-{(2-hydroxynaphthalene-1-yl) methylidene] amino} ethyl} naphthalene-2-ol

Actual yield = 4.68

Theoretical yield = 6.79

% yield = 68.92%

(iii) Co-1-{(E)-[2-{(2-hydroxynaphthalene-1-yl) methylidene] amino} ethyl} naphthalene-2-ol

Actual yield = 4.80g

Theoretical yield = 6.19g

% yield = 78.12%

FT – IR Result

Table 1: IR Results

EDNA (cm^{-1})	NiEDNA (cm^{-1})	CoEDNA (cm^{-1})	Band Assignment
1614	1603	1618	V(C = N)
3679	3347	3753	V(O – H)
1204	1252	1252	V(C – O)
3049	–	3056	V(Ar – H)

Conductivity Result

Table 2: Conductivity of 1% solutions of the Complexes

Complex	Conductivity μscm^{-1}
NiEDNA	51.6
CoEDNA	71.6

Elemental Analysis of EDNA

Table 3: Elemental Analysis result for EDNA

Element	Calculated	Found
C	78.26	64.40
H	5.43	6.10
N	7.61	5.60

Inhibitory activities of the Schiff base and its complexes of nickel (II) and cobalt (II) ions.

Table 4: Inhibitory activities of complexes on selected pathogens

Complexes	Inhibitory zone diameters, IZD (in cm)				
	B. subtilis	*S. aureus*	*S. sp*	*E. coli*	*P.aeruginosa*
CoEDNA	2.5	2.0	1.6	1.8	1.0
NiEDNA	1.2	Nil	Nil	1.1	Nil
EDNA	1.2	Nil	Nil	1.2	Nil

Inhibitory activities of the Schiff base and its complexes of nickel (II) and cobalt (II) ions.

Table 5: Inhibitory activities of complexes on selected pathogens

Complexes	Inhibitory zone diameters, IZD (in cm)				
	B. subtilis	*S. aureus*	*S. sp*	*E. coli*	*P.aeruginosa*
CoEDNA	2.5	2.0	1.6	1.8	1.0
NiEDNA	1.2	Nil	Nil	1.1	Nil
EDNA	1.2	Nil	Nil	1.2	Nil

MIC (µg/ml) result

Table 6: MIC (µg/ml) of the complexes against Gram-positive and Gram-negative bacteria.

Compound	*B. subtilis*	*S. aureus*	*Str. sp*	*E. coli*	*P.aeruginosa*
CoEDNA	20	500	500	500	500
NiEDNA	500	-	-	500	-
EDNA	500	-	-	500	-

Discussion

A Schiff base 1-{(E)-[2-{(2-hydroxynaphthalene-1-yl) methylidene] amino} ethyl} naphthalene-2-ol was derived from ethylenediamine and 2-hydoxy-1-naphthaldehyde and its complexes with Ni (II) and Co (II) ions were synthesized. The ligand is yellow in colour with a melting point of 310 °C, its nickel (II) complex is greenish-brown while its cobalt (II) complex is reddish-pink in colour. There is appearance of absorption at around 1600 cm^{-1} in the

IR spectra indicating the presence of the azomethine group (N=C), the complexes are characterized by molar absorptivity above 1000 $Lmol^{-1}cm^{-1}$ which is not seen in the ligand from which they were derived indicating modification in the electronic transition that led to stronger absorption in the UV-Visible region of the electromagnetic radiation.

The anti-microbial assay carried out on some selected microorganisms showed the cobalt complex inhibits growth in *B. subtilis, S. aureus, S.sp, E. coli,* and *P. aeruginosa* while both the nickel complex and the ligand show growth inhibition only on *B. subtilis,* and *E. coli.* The ligand and its nickel and cobalt (II) complexes are only soluble in dimethylsolphoxide (DMSO) and dimethylformamide (DMF) at high temperature which led to unsuccessful attempt at ^{1}H and ^{13}C NMR.

Conclusion

The spectroscopic data obtained shows the successful synthesis of the Schiff base 1-{(E)-[2-{(2-hydroxynaphthalene-1-yl) methylidene] amino} ethyl} naphthalene-2-ol. The differences in the molar absorptivity and the anti-microbial activities between the complexes and the ligand from which they were derive proved the successful modification of the Schiff base through formation of complexes.

REFERENCES

Alam M. S., Choi J. H., Lee Du (2012). Synthesis of novel Schiff base analogues of 4-Amino- 1,5-diamethyle-2-phenylpyrazol-3-one and their evaluation for anti-inflammatory activity. *Bioorganic medical chemistry*.Vol. 20. Pp.4103.

Arumugam AP, Guhanathan S. and Elango G, (2017). Co (II), Ni (II) and Cu (II) Complexes with Schiff Base Ligand: Syntheses, Characterization, Antimicrobial Studies and Molecular Docking Studies, SOJ Materials Science & Engineering


IUPAC, (the "Gold book") (1997). Compendium of chemical Terminology, 2nd ed. Online corrected version: (2006).


Neclakantan MA, Rusalraj F, Dhrmaraja J, Johnsonraajas, Jeygkumar T, Pillai ms. (2008). Specified Characterization, Cyclic voltammetry, Morphology, biological activities and DNA Clearing studies of Amino Schiff Base's Metal complexes. *Spectrochum Acta.* Vol. 71A. Pp. 1599-609.


Obasi Nnamdi Lawrence, Uchechukwu Susan Oruma, Ibrahim Abdulrazak Alswaidan, Ponnadurai Ramasami, Chigozie Julius Ezeorah and Alfred Ezinna Ochonogor (2017). Synthesis, Characterization and anti-bacterial Studies of N-(Benzothiazol-2-yl)-4-Chlorobenzenesulphonamide and its Neodymium(III) and Thallium(III) Complexes. *Molecules*. Vol. 22(2). Pp. 152.

Selma Bal and Sedat Salih Bal, (2014). Cobalt(II) and Manganese(II) Complexes of Novel Schiff Bases, Synthesis, Charcterization, and Thermal, Antimicrobial, Electronic, and Catalytic Features. Hindawi Publishing Corporation, Advances in Chemistry, Volume 2014, Article ID 506851.

Wang RM, Maojj, Song JF, Huo Cx, He YF.(2007). Anti-oxidant activity of Bovine serum albumin binding amino acid schiff base's metal complexes. *Chinese chemical letter.* Vol.18:14. Pp.16-8.









EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]

Agilent Technologies

Sample ID:EDNA
Sample Scans:30
Background Scans:16
Resolution:8
System Status:Good
Method Name:Transmittance Method
User:Admin
Date/Time:2018-04-24T17:07:00.437+01:00
Range:4000 - 650
Apodization:Happ-Genzel
File Location:C:\Program Files\Agilent\MicroLab PC\Results\EDNA_2018-04-24T17-07-00.a2r




Sample ID:NiEDNA
Sample Scans:30
Background Scans:16
Resolution:8
System Status:Good
Method Name:Transmittance Method
User:Admin
Date/Time:2018-04-24T17:01:31.817+01:00
Range:4000 - 650
Apodization:Happ-Genzel
File Location:C:\Program Files\Agilent\MicroLab PC\Results\NiEDNA_2018-04-24T17-01-31.a2r



EDITOR-IN-CHIEF
Prof. A.S. Hossein [Saskatchewan]

Sample ID:CoEDNA
Sample Scans:30
Background Scans:16
Resolution:8
System Status:Good
Method Name:Transmittance Method
User:Admin
Date/Time:2018-04-24T17:04:12.454+01:00
Range:4000 - 650
Apodization:Happ-Genzel
File Location:C:\Program Files\Agilent\MicroLab PC\Results\CoEDNA_2018-04-24T17-04-12.a2r



EDNA: for this sample we couldn't have 1H and 13C Spectra, for reasons of solubility. Even with hot DMSO, it dissolve and instantly return to build a precipitate.




Figure 13. IR Spectra of compound **EDNA**.

- Elemental analysis relating to compound **EDNA**

Anal. Calcd (%) for $C_{33}H_{26}ON_2$: C 78.26, H 5.43, N 7.61. Found: C 64.4, H 6.1, N 5.6.

The sample does not have a stable weigh.

